

Patient: \_\_\_\_\_

Date: \_\_\_\_\_

Notes:

Cephalic Flow

\_\_\_\_\_ ml/min

Basilic Flow

\_\_\_\_\_ ml/min

Basilic

Cephalic

Elbow  
Joint

endoAVF  
Select One

Radial - Radial

Ulnar - Ulnar

### KEY

Cannulation Zone



Description:

This diagram represents the flow rates and recommendations for cannulation of the endoAVF.  
If you have any questions, please call the number below:

Name \_\_\_\_\_

Number \_\_\_\_\_

The WavelinQ™ EndoAVF System is indicated for the creation of an arteriovenous fistula (AVF) using concomitant ulnar artery and ulnar vein or concomitant radial artery and radial vein in patients with minimum artery and vein diameters of 2.0 mm at the fistula creation site who have chronic kidney disease and need hemodialysis. **Please consult product labels and instructions for use for indications, contraindications, hazards, warnings and precautions.**

[bd.com/wavelinq](http://bd.com/wavelinq) BD, Tempe, AZ, USA, 1 800 321 4254

BD, the BD logo, and WavelinQ are trademarks of Becton, Dickinson and Company or its affiliates. © 2022 BD. All rights reserved.  
© 2022 Illustration by Mike Austin. BD-15292v2

